

Cercartetus concinnus (Diprotodontia: Burramyidae)

JAMIE M. HARRIS

School of Environmental Science and Management, Southern Cross University, Lismore, New South Wales, 2480, Australia; jharrill@scu.edu.au

Abstract: *Cercartetus concinnus* (Gould, 1845) is a burramyid commonly called the western pygmy-possum. It is 1 of 4 species in the genus *Cercartetus*, which together with *Burramys parvus* form the marsupial family Burramyidae. *C. concinnus* is endemic to southern mainland Australia. Its habitat is heathlands, shrublands, and dry forests in semiarid zone areas and it feeds on a range of foods, particularly nectar and pollen. This species is notable for short-term hibernation and its role as a pollinator of plants. It is considered secure throughout most of its range except for southwestern New South Wales, where it is classified as endangered. DOI: 10.1644/831.1.

Key words: Australia, Burramyidae, hibernator, marsupial, pollinator, western pygmy-possum

Published 27 May 2009 by the American Society of Mammalogists
Synonymies completed 1 May 2008



Cercartetus Gloger, 1841

- Cercartetus* Gloger, 1841:85. Type species *Phalangista nana* Desmarest, 1817, by monotypy.
- Dromicia* Gray, 1841:401. Type species *Phalangista nana* Desmarest, 1817, by monotypy.
- Burramys* Broom, 1896:563. Type species *Burramys parvus* Broom, 1896, by monotypy.
- Eudromicia* Mjöberg, 1916:13. Type species *Eudromicia macrura* Mjöberg, 1916, by subsequent designation (Matschie 1916:260).
- Dromiciola* Matschie, 1916:260. Type species *Dromicia lepida* Thomas, 1888, by original designation.
- Dromiciella* Matschie, 1916:260. Type species *Dromicia concinna* Gould, 1845, by original designation.

CONTEXT AND CONTENT. Order Diprotodontia, suborder Phalangeriformes, superfamily Phalangoidea, family Burramyidae (Kirsch 1968). The family Burramyidae consists of 2 genera: *Burramys* and *Cercartetus*. Members of *Cercartetus* are *C. caudatus*, *C. concinnus*, *C. lepidus*, and *C. nanus* (Harris 2008; van Dyck and Strahan 2008).

A rudimentary key to the extant burramyids adapted from descriptions in Flannery (1994) follows:

- 1. Ventral fur pure white *C. concinnus*
Ventral fur more gray than white 2
- 2. Adult body weight ≤9 g *C. lepidus*
Adult body weight substantially >9 g 3
- 3. Tail about the same length as body and <100 mm *C. nanus*
Tail >1.2 times length of body and >140 mm 4

- 4. P3 longer than each molar, grooved and serrated *B. parvus*
P3 smaller than molars, not grooved or serrated *C. caudatus*

Cercartetus concinnus (Gould, 1845)
Western Pygmy-possum

Dromicia concinna Gould, 1845:2. Type locality “Western Australia;” restricted to “Swan River” by Thomas (1922:127).



Fig. 1.—An adult *Cercartetus concinnus* from Nepean Bay Conservation Park, Kangaroo Island, South Australia. Used with permission of the photographer A. C. Robinson.

Phalangista (Dromicia) concinna: Waterhouse, 1846:314. Name combination.

Phalangista (Dromicia) neillii Waterhouse, 1846:315. Type locality “King George’s Sound, Western Australia.”

Dromicia (Dromiciella) concinna: Glauert, 1933:24. Name combination.

Cercartetus concinnus: Iredale and Troughton, 1934:22. First use of current name combination and correction for gender.

CONTEXT AND CONTENT. Context as for genus. Two subspecies (listed below) have been previously recognized (Flannery 1994; McKay 1988). However, as discussed in the Nomenclatural Notes, this is unwarranted, and accordingly I do not recognize them as valid.

C. concinnus concinnus Gould, 1845:2. See above.

C. concinnus minor Wakefield, 1963:100. Type locality “Nurcong [approximately 16 km] northwest of Natimuk,” Victoria.

NOMENCLATURE NOTES. The genus name *Dromicia* (Gray 1841) was used in the 1st description of the subject species (Gould 1845). An alternate genus, *Dromiciella* (Matschie 1916), was later proposed for *D. concinna*, although this was largely ignored. Iredale and Troughton (1934) noted that *Cercartetus* (Gloger 1841) antedated *Dromicia* by several months. They advanced *Cercartetus concinnus* to supersede *D. concinna* and this included a slight change in the epithet to accord with the new genus.

Two subspecies of *C. concinnus* were originally proposed based on slight differences in size and coloration but principally because of an apparent disjunct distribution: *C. c. concinnus* in southwestern Western Australia and *C. c. minor* in South Australia, western Victoria, and western New South Wales (Wakefield 1963). However, the subspecific arrangement was later questioned (Wakefield 1970) because of both modern and fossil records from approximately the middle of the distributional gap (Bolam 1923; Lundelius 1957), and lack of significant differences in body and skull measurements (Ryan 1963). Consequently, for some years the status of the 2 subspecies was a matter of debate (Flannery 1994; McKay 1988; Smith 1995), but recent genetic studies provide reasonably convincing evidence that there are in fact no subspecies (Osborne and Christidis 2002; Pestell et al. 2008).

It has occasionally been presumed that *Cercaertus* (Burmeister 1837) was a misspelling or synonym of *Cercartetus* (e.g., Cook 1963; Grzimek 1975; Ryan 1963; Sharman 1963; Simpson 1945). However, *Cercaertus* is a junior synonym of *Trichosurus* (brushtail possums) and not of *Cercartetus* (Harris 2006; Wakefield 1963). *Cercartetus* is a reference to the prehensile tail (from the Greek *kerkos*) and *concinnus* is derived from the Latin word for elegant or neat (Flannery 1994; Strahan 1981).

Vernacular names used include “beautiful pigmy-phalanger” (Waterhouse 1846), “Neill’s phalanger” (Gerrard 1862;

Waterhouse 1846), “beautiful *Dromicia*” (Gould 1863), “King George’s Sound *Dromicia*” (Krefft 1871), “lesser dormouse-phalanger” (Thomas 1888), “western dormouse-opossum” (Ogilby 1892), “western dormouse-phalanger” (Lydekker 1896), “elegant dormouse possum” (Wood Jones 1924), “western pigmy possum” (Iredale and Troughton 1934), and “thin-tailed dormouse possum” or “southwestern pigmy phalanger” (Grzimek 1975). Aboriginal names for *C. concinnus* include “Man-dur-da” (Gould 1863; Whittell 1954) or “Mundarda” (Brooker 1988; Wakefield 1963), “Jāā-jat” (Waterhouse 1846), “Nyeranit” (Abbott 2001; Shorridge 1909), and “Wirappi” (Tunbridge 1991).

DIAGNOSIS

Cercartetus concinnus (Fig. 1) is distinguished from other pigmy-possums by its size, coloration, length of tail, and dentition. This species has pure white underparts (including chin and lower jaw), and the line of demarcation from the bright cinnamon-gray dorsal fur is sharply marked. All other pigmy-possum species have gray-based belly fur.

GENERAL CHARACTERS

Cercartetus concinnus has a delicate skull with short, pointed rostrum, slender zygomatic arch, and broad squamosals (Fig. 2; Rosenberg and Richardson 1995; Turner and McKay 1989). The dentition is diprotodont and the 3 molars present have simplified crowns. It possesses long vibrissae, large, black, forward-directed eyes, and a dark patch in front of the eyes, but this is indistinct compared with that of some other pigmy-possum species. The ears are large, oval-shaped, thin, and membranous (Fig. 1). The tail is finely scaled, long, slender, prehensile, and cylindrical, except for the base, which is occasionally incrassated (Casanova 1958). On the digits of the forefeet and hind feet they have expanded apical pads (Fig. 3; Wood Jones 1924) that have eccrine sweat glands (Green 1963; Rosenberg and Rose 1999). The 2nd and 3rd digits on the hind feet are syndactylous, and the hallux is opposable and clawless.

The sexes are very similar in size and color (Gould 1863). Adult body weight is 8–21 g (\bar{X} = 13 g—Carthew et al. 2008). Ranges and means (mm) for standard external measurements from 4 male specimens in the Australian Museum are: length of head and body (57–72; 66); length of tail (77–87; 81); length of hind foot (11.0–13.0; 11.5); length of ear (13.0–15.0; 14.5—Flannery 1994). Further data of this type are available elsewhere (Carthew et al. 2008; Thomas 1888; Wakefield 1963; Waterhouse 1846; Wood Jones 1924, 1925).

DISTRIBUTION

Cercartetus concinnus (Fig. 4) occurs in southwestern Western Australia (Dell and How 1985; Glauert 1933; How






Fig. 4.—Geographic distribution of *Cercartetus concinnus*. Adapted from van Dyck and Strahan (2008) and Pearson et al. (1999).

these sites are late Pleistocene to Holocene (see Lundelius 1983).

FORM AND FUNCTION

Dental formula is $i\ 3/2$, $c\ 1/0$, $p\ 3/3$, $m\ 3/3$, total 36. Technical descriptions and measurements of the dentition, skull, and face musculature are available in the literature (Osborne and Christidis 2002; Rosenberg and Richardson 1995; Ryan 1963; Thomas 1888; Wakefield 1963; Waterhouse 1846). The tongue is broad, long (10–12 mm), tapers to a blunt point, and the rostral half reportedly lacks a brushlike structure (Rosenberg and Richardson 1995). All burramyids have a simple gut structure and a well-developed cecum (Turner and McKay 1989). The kidneys of *Cercartetus concinnus* have a high maximal urinary concentrating capacity (6102 mOsm/kg H₂O—Slaven and Richardson 1988).

The female's pouch is well developed and typical of diprotodont marsupials (type 5—Russell 1982). It opens anteriorly and contains 6 teats. Males have a nonpendulous scrotum (Smith 1984) and carrot-shaped prostrate (Ward 1990). Measurements of testis length (4.3–4.5 mm) and width (3.2–3.7 mm) are available from museum specimens ($n = 2$ —Woolley and Vanderveen 2002). Both male and female *C. concinnus* possess paracloacal glands (Green 1963; Smith 1984)

←

Fig. 3.—Palmar surface of the manus and pes of *Cercartetus concinnus*. Reproduced from Wood Jones (1924).

and have reproductive tracts that are similar to those of other species of *Cercartetus* (Turner and McKay 1989; Ward 1990).

All burramyids investigated in some detail have the ability to hibernate (i.e., undergo prolonged bouts of torpor—Geiser 1985, 1987). In *C. concinnus*, the longest torpor bout recorded is 11 days (Wakefield 1970). Observations under laboratory conditions were 4 days at an ambient temperature of 12°C and 8 days at an ambient temperature of 8°C (Geiser 1987). Mean body temperature at normothermia was 34.4°C \pm 0.5°C *SD*; during torpor body temperature dropped to as low as 4.7°C (Geiser 1987; see also Zosky and Larcombe 2003). Oxygen consumption (a measure of metabolic rate) of torpid individuals was 0.046 L kg⁻¹ h⁻¹ \pm 0.02 *SD* and increased about 200-fold during normal activity (Geiser 1987; see also Zosky and Larcombe 2003). Torpid individuals may have periods of apnea lasting for about 5 min at an ambient temperature of 10°C (Geiser 1987). Average heart rate of *C. concinnus* is 18.0 beats/min \pm 1.1 *SD* during torpor and 20.1 \pm 2.1 beats/min during eupnea (Zosky and Larcombe 2003). Arousal rates are rapid (over a 10-min period these averaged 0.79°C/min—Geiser 1987). Functionally, torpor allows *C. concinnus* to reduce overall energy expenditure and increase fasting endurance during poor weather, lower temperatures, and during periods of food shortage (Fig. 5; Bennett and Lumsden 1995; Pearson et al. 1999; Turner and McKay 1989; Wakefield 1970).

ONTOGENY AND REPRODUCTION

It has been implied that embryonic diapause (or delayed implantation) occurs in *Cercartetus concinnus* (Kean 1967; Marshall 1967; Renfree 1981, 1993; Renfree and Calaby 1981; Sharman 1963; Ward 1990). This is based on anecdotal observations of captive females that produced a 2nd litter in the absence of a male (see Bowley 1939; Casanova 1958; Hartman 1940). Further investigations of this phenomenon found that the blastocysts continued to grow during lactation (Clark 1967). Although knowledge of reproduction in *C. concinnus* is incomplete, examination of available data suggests that diapause, *sensu stricto*, does not actually occur (see also Renfree 1978; Turner and McKay 1989; Tyndale-Biscoe 1973; Ward 1990). However, only about 2 days are needed between the weaning of one litter and commencement of suckling of the next; in that short period the mammary section changes from milk to colostrum and the enlarged teats regress to a size small enough to be taken into the mouth of a neonate (Clark 1967; Flannery 1994; Smith 1995).

Females are thought to reach sexual maturity at 12–15 months (Tyndale-Biscoe and Renfree 1987). The reproductive rate is high because breeding may occur throughout the year and females can rear 2 or 3 litters in close succession (Clark 1967; New South Wales National Parks and Wildlife Service 2001; Ward 1990). Records of lactating females and



Fig. 5.—Torpid *Cercartetus concinnus*. Note that torpid individuals usually sit on their tails with their backs exposed upward and curled as seen in the photograph. However, the photograph displays a possum on its side for ease of visualization. Used with permission of the photographer J. Turner.

juveniles are concentrated in the spring months and few in the winter months (Ward 1990). Females typically ovulate more eggs and carry more embryos than there are teats, so number of teats limits litter size to 6 (Ward 1990, 1998). There are a number of observations of 6 young (Bowley 1939; Casanova 1958; Glauert 1950) and some records of only 4 young (Jones 1954; Pearson et al. 1999; Ryan 1963). In one field study, the mean number of pouch young was 4.6 \pm 1.0 *SD* (n = 21 litters—Ward 1990, 1998).

Newborns have a head length of about 2.9 mm (Clark 1967). The young leave the pouch at around 25 days before the eyes open (Bowley 1939; Clark 1967) but remain in a nest (Bowley 1939; Casanova 1958). At this stage, the young are only sparsely furred and are dependent on the mother. The young are suckled until they reach about 50 days (Clark 1967).

ECOLOGY

Population characteristics.—There is some evidence to indicate that populations fluctuate widely in numbers, depending on climatic conditions and the availability of resources (Carthew 2004; Paton 1999; Pearson et al. 1999; Pestell and Petit 2007b; Rose 1995). At Queen Victoria Springs, Western Australia, lack of recaptures between trapping sessions suggested that *Cercartetus concinnus* is a transient in its habitat, either dispersing through it in search of preferred habitat or visiting the area on a nomadic basis in search of food resources or mates (Pearson et al. 1999). To



Fig. 6.—Habitat of *Cercartetus concinnus* in Newland Head Conservation Park, South Australia. This site is coastal mallee heath dominated by *Banksia marginata*, *B. ornata*, *Callistemon rugulosus*, and a mix of eucalypt species, including *Eucalyptus baxteri*, *E. cosmophylla*, and *E. diversifolia*. Used with permission of the photographer S. Carthew.

maximize capture success, surveys of *C. concinnus* (using pitfall traps or nest boxes) should be planned to coincide with flowering of proteaceous and myrtaceous understory species and avoid periods of inclement weather when the species is likely to be torpid (New South Wales National Parks and Wildlife Service 2001).

Space use.—*Cercartetus concinnus* occurs in areas with high temperatures and low rainfall (Ryan 1963) and is essentially a semiarid-zone species (Brown 1974). Its habitat is generally stated to be mallee woodlands, heathlands, shrublands, and dry forests with a heathy or structurally dense understory of callistemons and melaleucas (family Myrtaceae) and banksias, grevilleas, and hakeas (family Proteaceae—Fig. 6; Bennett and Lumsden 1995; Carthew et al. 2008; Flannery 1994; Kemp and Carthew 2004; Menkhorst and Beardsell 1982; Misso 1997; Wakefield 1963; Ward 1992). At Mallee Cliffs National Park, New South Wales, *C. concinnus* was trapped in unusual habitat—a belah (*Casuarina pauper*) patch in a mixed woodland with an understory of cottony saltbush (*Chenopodium curvispicatum*), bitter saltbush (*Atriplex stipitata*), ruby saltbush (*Enchylaena tomentosa*), and Mueller's daisy bush (*Olearia muelleri*—Fig. 7; New South Wales National Parks and Wildlife Service 2001). Also in New South Wales, at Boree Plains Station, the species was recorded from mallee spinifex shrubland with an understory of *Leptospermum coriaceum* (see New South Wales National Parks and Wildlife Service 2001).

Cercartetus concinnus shelters in a variety of locations, including tree hollows, under loose pieces of bark, in cavities excavated by fire, under the fronds of *Xanthorrhoea*, between dead *Banksia* inflorescences, in abandoned birds' nests (e.g.,



Fig. 7.—Habitat of *Cercartetus concinnus* in Mallee Cliffs National Park, New South Wales. Used with permission of the photographer J. Smith.

grey-crowned babbler [*Pomatostomus temporalis*] and Zebra Finch [*Taeniopygia guttata*], among leaves on the ground, at the base of trees, and in hollow stumps (Aitken 1954; Casanova 1958; Flannery 1994; Gould 1863; Kemp and Carthew 2004; Lydekker 1896; Misso 1997; Pestell and Petit 2007a, 2007b; Ryan 1963; Shortridge 1909; Tulloch 2004; Wakefield 1963; Waterhouse 1846; Wood Jones 1924). Different types of shelter sites may be used for different purposes, such as breeding and avoidance of predators (Misso 1997; Tulloch 2004).

Little information exists on the movements and size and spacing of home ranges of this species. However, in southeastern South Australia, the mean distance moved by individuals between captures ranged from 24 m to 60 m, and the maximum distance recorded between captures was 195 m (Horner 1994; not seen, cited in New South Wales National Parks and Wildlife Service 2001). Males tended to move further than females. In the Middleback Ranges in South Australia, mean distances travelled were 56 m for males and 50 m for females and the maximum recorded distance was 195 m (S. Carthew, in litt.).

Diet.—*Cercartetus concinnus* feeds on a range of foods including nectar, pollen, and insects (Cadzow and Carthew 2004; Glauert 1950; Horner 1994; Pestell and Petit 2007b; Wakefield 1963; Waterhouse 1846; Whittell 1954). Nectar and pollen are probably the primary dietary components, and sources include species in the genera *Banksia*, *Eucalyptus*, and *Melaleuca* (Cadzow and Carthew 2004; Pestell and Petit 2007b). *C. concinnus* is a regular and nondestructive flower visitor and may play an important part in the pollination of its food plants (Carthew et al. 2008; Pestell and Petit 2007b; Turner 1982; Turner and McKay 1989). In captivity, *C. concinnus* may accept soft fruit, apples, cakes, rose leaves, jam, honey, cow's milk, baby food, honey–water mixture, custard, freshly collected *Banksia* pollen, and mealworms (*Tenebrio molitor*—Casanova 1958; Geiser 1987; Glauert 1950; Landwehr et al. 1990; Wakefield 1970; Wood Jones 1924; Wooller

et al. 1999; Zosky and Larcombe 2003). A laboratory study of sugar preferences in *C. concinnus* found no clear preference for any one particular sugar type (Landwehr et al. 1990). Another laboratory study determined daily nitrogen requirements to be about $0.146 \text{ g N/kg}^{0.75} \pm 0.086 \text{ SD}$ (Wooller et al. 1999).

Diseases and parasites.—Fleas recorded as parasitic on *Cercartetus concinnus* include *Acanthopsylla scintilla scintilla* and *A. scintilla tasmanica* (Dunnet and Mardon 1974; Holland 1971). Astigmatic mite records are for *Listrophoroides queenslandicus* and *Murichirus notomys* (Domrow 1992). Hepatoma (a type of cancer originating in the liver) is known from captive *C. concinnus*, although the etiology of the condition remains unknown and it is unclear whether this is a common cause of death in this species (Canfield et al. 1990; Hopkins et al. 1984; Turner and McKay 1989).

Interspecific interactions.—The domestic cat (*Felis catus*) is a known predator of *Cercartetus concinnus* (Loaring 1960; Wood Jones 1924). In addition, the introduced red fox (*Vulpes vulpes*) and a range of native Australian carnivorous marsupials (e.g., quolls [*Dasyurus*]), snakes, and owls probably prey regularly on *C. concinnus*. Sympatry with *C. lepidus* occurs in a number of areas including Ngarkat Conservation Park in South Australia (Paton 1999); the Big Desert, Victoria (Bennett et al. 1989; Robertson et al. 1989; Ward 1992); and Jilpanger Flora and Fauna Reserve, Victoria (Conole 1996). It is not yet clear how these ecologically similar congeners partition the habitat and other resources such as food and nesting sites (Bennett et al. 1989; Cadzow and Carthew 2004; Robertson et al. 1989).

Miscellaneous.—The species can be maintained readily in captivity (Aitken 1954; Brooker 1988; Casanova 1958; Fairfax 1982; Geiser 1987; Glauert 1950; Hopkins et al. 1984; Landwehr et al. 1990; Wakefield 1970; Waterhouse 1846; Wood Jones 1924; Wooller et al. 1999; Zosky and Larcombe 2003). However, individuals have a tendency to overeat and become obese (Fairfax 1982; Geiser 1987; George 1990; Landwehr et al. 1990; Wakefield 1970). Although breeding is known to have occurred at Perth Zoo, such instances have been irregular (George 1990). The author also is aware that about 20 years ago, a captive breeding population was maintained at Murdoch University in Perth, Western Australia, although this no longer exists.

BEHAVIOR

Cercartetus concinnus is scansorial, nocturnally active, and usually agile but individuals may be placid when handled (Carthew et al. 2008; Casanova 1958; Smith 1995; Wood Jones 1924). The forepaws are used to hold food or to hold open floral parts for better access to nectar and pollen (Casanova 1958; Turner and McKay 1989; Wood Jones 1924). The prehensile tail is used to cling to branches or grasp objects such as nesting materials (Bolam 1923; Casanova 1958). During torpor, *C. concinnus* typically coils tightly into

a ball with the ears folded over the eyes and the tail spiraled upon the stomach (Fig. 5; Bolam 1923; Casanova 1958; Pearson et al. 1999; Wakefield 1970). In hot weather, *C. concinnus* has been observed panting (Casanova 1958). Extensive self-grooming is undertaken after feeding bouts (Casanova 1958). Its vocalization is described as a rapid chattering “chi-chi-chi” (Casanova 1958).

Antipredator behavior in burramyids is thought to include anachoresis (avoidance by concealment in a retreat, such as a burrow, that precludes an attack), specific activity patterns (such as adjustment of foraging activity to minimize predation risk), vigilance (a general wariness), withdrawal (retreat to hollow or mother’s pouch when alarmed), retaliation (biting or intimidatory displays including hissing), and flight (fleeing from the predator—Coulson 1996).

GENETICS

Cercartetus concinnus has XY/XX sex determination and a diploid number (2n) of 14 chromosomes (Martin and Hayman 1967; McKay 1984; Rofe and Hayman 1985). It has been suggested that the population of *C. concinnus* in New South Wales may be genetically distinct from nearby Victorian populations (New South Wales National Parks and Wildlife Service 2001). However, a number of recent molecular analyses (mitochondrial DNA [mtDNA] *ND2* gene [Osborne and Christidis 2002] and mtDNA *ND4* gene [Pestell et al. 2008]) found <1% sequence divergence among samples from widely distributed populations. This casts doubt on any special genetic significance of the New South Wales population, and also that *C. concinnus* represents a single monophyletic group across its range (see also subspecific dichotomy in Nomenclatural Notes above). *C. concinnus* has a sister relationship with *C. nanus* (Osborne and Christidis 2002), and probably diverged in the late Miocene about 8 million years before present (Pestell et al. 2008).

CONSERVATION

The species is listed as “Least concern, Lower risk” by the International Union for the Conservation of Nature and Natural Resources (2006), and is not listed as a threatened species federally in Australia or in state-based legislation in Western Australia, South Australia, or Victoria (Tulloch 2004). However, in New South Wales, *Cercartetus concinnus* is listed as endangered (Kavanagh 2004; New South Wales National Parks and Wildlife Service 2001; New South Wales Scientific Committee 1997). In this state, *C. concinnus* is threatened by vegetation clearing, the reduction of food sources by livestock overgrazing, inappropriate fire regimes, and introduced predators (i.e., feral cats and foxes—New South Wales National Parks and Wildlife Service 2001; New South Wales Scientific Committee 1997; Rose 1995; Tulloch

2004). Potentially, the harvesting of nectar and pollen by commercial honeybees also could have a detrimental impact on populations of *C. concinnus*, although this has been little studied (Paton 1999).

A few informal assessments of its conservation status in specific areas are available in the literature, and for its range in Western Australia these include that it is “common” (Baynes 1970; Glauert 1933) or “stable” (Burbidge and McKenzie 1989). However, in the Western Australian wheat-belt *C. concinnus* is reportedly “rare” (Kitchener et al. 1980). In most of its South Australian range, *C. concinnus* is considered “common” (Aitken 1983; Carthew 2004; Robinson and Armstrong 1999; Watts 1990; Watts and Ling 1985), except for Brookfield Conservation Park where it is “rare” (Goonan et al. 1993). In the mallee of northwestern Victoria, *C. concinnus* is “common” to “uncommon” (Bennett and Lumsden 1995; Bennett et al. 1989; Robertson et al. 1989; van der Ree et al. 2004). However, a southward contraction at the northern end of its geographic range in Victoria has been noted (Bennett and Lumsden 1995).

ACKNOWLEDGMENTS

I am grateful to S. Carthew, K. Munro, A. C. Robinson, J. Smith, J. Turner, and S. van Dyck for assistance with the figures and to T. du Bois, I. Abbott, and 1 anonymous reviewer for helpful comments to improve this account.

LITERATURE CITED

- ABBOTT, I. 2001. Aboriginal names of mammal species in south-west Western Australia. *CALM Science* 3(4):433–486.
- AITKEN, P. F. 1983. Mammals. Pp. 127–134 in *The natural history of the south-east* (M. J. Tyler, J. K. Ling, and J. W. Homes, eds.). Royal Society of South Australia, Adelaide, Australia.
- AITKEN, R. 1954. The recent increase of the rarer native mammals. VI. Nyabing-Pingrup. *Western Australian Naturalist* 4:138–139.
- ANONYMOUS. 1914. Remarks by exhibitors of specimens. *Victorian Naturalist* 31:50–52.
- ARCHER, M. 1974. New information about the Quaternary distribution of the thylacine (Marsupialia: Thylacinidae) in Australia. *Journal of the Royal Society of Western Australia* 57:43–50.
- ARCHER, M., AND A. BAYNES. 1972. Prehistoric mammal faunas from two small caves in the extreme south-west of Western Australia. *Journal of the Royal Society of Western Australia* 55: 80–89.
- BALME, J., D. MERRILEES, AND J. K. PORTER. 1978. Late Quaternary mammal remains, spanning about 30,000 years, from excavations in Devil's Lair, Western Australia. *Journal of the Royal Society of Western Australia* 61:33–65.
- BAYNES, A. 1970. Catching pygmy possums. *Western Australian Naturalist* 11:189–190.
- BAYNES, A. 1987. The original mammal fauna of the Nullarbor and southern peripheral regions: evidence from skeletal remains in superficial cave deposits. Pp. 139–152, 400–401 in *A biological survey of the Nullarbor region of South and Western Australia in 1984* (N. L. McKenzie and A. C. Robinson, eds.). South Australian Department of Environment and Planning, Adelaide, Australia.
- BAYNES, A., D. MERRILEES, AND J. K. PORTER. 1975. Mammal remains from the upper levels of a late Pleistocene deposit in Devil's Lair, Western Australia. *Journal of the Royal Society of Western Australia* 58:97–126.
- BENNETT, A. F., AND L. F. LUMSDEN. 1995. Western pygmy-possum *Cercartetus concinnus*. Pp. 97–98 in *Mammals of Victoria—distribution, ecology and conservation* (P. W. Menkhorst, ed.). Oxford University Press, Melbourne, Australia.
- BENNETT, A. F., L. F. LUMSDEN, AND P. W. MENKHORST. 1989. Mammals of the mallee, southeastern Australia. Pp. 191–220 in *Mediterranean landscapes in Australia, mallee ecosystems and their management* (J. C. Noble and R. A. Bradstock, eds.). CSIRO Publishing, Melbourne, Australia.
- BOLAM, A. G. 1923. *The trans-Australian wonderland*. Modern Printing Company, Melbourne, Australia.
- BOWLEY, E. A. 1939. Delayed fertilization in *Dromicia*. *Journal of Mammalogy* 20:449.
- BROOKER, B. 1988. *Mundarda*. University of Western Australia Press, Nedlands, Australia.
- BROOM, R. 1896. On a small fossil marsupial with large grooved premolars. *Proceedings of the Linnean Society of New South Wales* 10:563–567.
- BROWN, G. D. 1974. The biology of marsupials of the Australian arid zone. *Australian Mammalogy* 1:269–288.
- BURBIDGE, A. A., AND N. L. MCKENZIE. 1989. Patterns in the modern decline of Western Australia's vertebrate fauna: causes and conservation implications. *Biological Conservation* 50:143–198.
- BURMEISTER, C. H. C. 1837. *Phalangista*. *Handbuch der Naturgeschichte* 2:814.
- CADZOW, B. R., AND S. M. CARTHEW. 2004. The importance of two species of *Banksia* in the diet of the western pygmy-possum *Cercartetus concinnus* and the little pygmy-possum *Cercartetus lepidus* in South Australia. Pp. 246–253 in *The biology of Australian possums and gliders* (R. L. Goldingay and S. M. Jackson, eds.). Surrey Beatty and Sons, Chipping Norton, Australia.
- CANFIELD, P. J., W. J. HARTLEY, AND G. L. REDDACLIFF. 1990. Spontaneous proliferations in Australian marsupials—a survey and review. I. Macropods, koalas, wombats, possums, and gliders. *Journal of Comparative Pathology* 103:135–146.
- CARTHEW, S. M. 2004. Distribution and conservation status of possums and gliders in South Australia. Pp. 63–70 in *The biology of Australian possums and gliders* (R. L. Goldingay and S. M. Jackson, eds.). Surrey Beatty and Sons, Chipping Norton, Australia.
- CARTHEW, S. M., B. R. CADZOW, AND J. N. FOULKES. 2008. Western pygmy-possum *Cercartetus concinnus*. Pp. 215–217 in *The mammals of Australia* (S. van Dyck and R. Strahan, eds.). 3rd ed. Reed New Holland, Chatswood, Australia.
- CASANOVA, J. 1958. The dormouse or pygmy possum. *Walkabout* 24: 30–31.
- CLARK, M. J. 1967. Pregnancy in the lactating pigmy possum, *Cercartetus concinnus*. *Australian Journal of Zoology* 15:673–684.
- COCKBURN, A., M. FLEMING, AND J. WAINER. 1979. The comparative effectiveness of drift fence pitfall trapping and conventional cage trapping of vertebrates in the Big Desert, north-western Victoria. *Victorian Naturalist* 96:92–95.
- CONOLE, L. E. 1996. The little pygmy-possum *Cercartetus lepidus*; an addition to the fauna of south-west Victoria. *Victorian Naturalist* 113:20–24.
- COOK, D. L. 1963. The fossil vertebrate fauna of Strongs' Cave, Boranup, Western Australia. *Western Australian Naturalist* 8: 153–162.
- COULSON, G. 1996. Anti-predator behaviour in marsupials. Pp. 158–186 in *Comparison of marsupial and placental behaviour* (D. B. Croft and U. Gansloßer, eds.). Filander Verlag, Fürth, Germany.
- DELL, J., AND R. A. HOW. 1985. The biological survey of the eastern goldfields of Western Australia. Part 3. Jackson–Kalgoorlie study area. IV. Vertebrate fauna. *Records of the Western Australian Museum Supplement* 23:39–66.
- DESMAREST, A. G. 1817. *Nouveau dictionnaire d'histoire naturelle*. Deterville, Paris, France 25:477.
- DOMROW, R. 1992. Acari Astigmata (excluding feather mites) parasitic on Australian vertebrates: an annotated checklist, keys and bibliography. *Invertebrate Taxonomy* 6:1459–1606.

- DORTCH, C. E., AND D. MERRILEES. 1971. A salvage excavation in Devil's Lair, Western Australia. *Journal of the Royal Society of Western Australia* 54:103–113.
- DUNNET, G. M., AND D. K. MARDON. 1974. A monograph of Australian fleas (Siphonaptera). *Australian Journal of Zoology, Supplementary Series* 30:1–273.
- FAIRFAX, R. A. 1982. Notes on the western pygmy-possum *Cercartetus concinnus* in captivity. P. 87 in *The management of Australian mammals in captivity* (D. D. Evans, ed.). Zoological Board of Victoria, Melbourne, Australia.
- FLANNERY, T. 1994. *Possums of the world: a monograph of the Phalangeroidea*. GEO Productions, Chatswood, Australia.
- GEISER, F. 1985. Hibernation in pygmy possums (Marsupialia: Burramyidae). *Comparative Biochemistry and Physiology, A. Comparative Physiology* 81:459–463.
- GEISER, F. 1987. Hibernation and daily torpor in two pygmy possums (*Cercartetus* spp., Marsupialia). *Physiological Zoology* 60:93–102.
- GEORGE, G. G. 1990. Monotreme and marsupial breeding programs in Australian zoos. *Australian Journal of Zoology* 37:181–205.
- GERRARD, E. 1862. *Catalogue of the bones of Mammalia in the collection of the British Museum*. Printed by Order of the Museum Trustees, London, United Kingdom.
- GLAUERT, L. 1933. The distribution of the marsupials in Western Australia. *Journal of the Royal Society of Western Australia* 19:17–32.
- GLAUERT, L. 1950. Observations on marsupials in captivity. *Western Australian Naturalist* 2(3):54–57.
- GLOGER, C. W. L. 1841. *Cercartetus*. Pp. 85 in *Gemeinnütziges Hand- und Hilfsbuch der Naturgeschichte*. Verlag von Aug. Schulz, Breslau, Germany (now Wrocław, Poland).
- GOONAN, P., A. L. MANN, AND T. P. MORLEY. 1993. Fauna of the Brookfield Conservation Park in the Murray mallee, South Australia. 2. Annotated checklist of the amphibians, reptiles and mammals. *South Australian Naturalist* 68:4–14.
- GOULD, J. 1845. *Dromicia concinna*. *Proceedings of the Zoological Society of London* 13:2.
- GOULD, J. 1863. *The mammals of Australia*. Reprint incorporating the 3 original volumes by modern notes by Joan M. Dixon. Macmillan 1977, South Melbourne, Australia.
- GRAY, J. E. 1841. Contributions towards the geographical description of the Mammalia of Australia, with notes on some recently discovered species. Pp. 397–414 (Appendix) in *Journal of two expeditions of discovery in northwest and Western Australia during the years 1837, 38 and 39* (G. Grey, ed.). T. and W. Boone, London, United Kingdom.
- GREEN, L. M. A. 1963. Distribution and comparative histology of cutaneous glands in certain marsupials. *Australian Journal of Zoology* 11:250–272.
- GRZIMEK, B. 1975. *Grzimek's animal life encyclopedia*. Vol. 10. Mammals I. Van Nostrand Reinhold, Melbourne, Australia.
- HARRIS, J. M. 2005. Annotated records of *Cercartetus* in the South Australian Naturalist. *South Australian Naturalist* 79(3–4):4–11.
- HARRIS, J. M. 2006. The discovery and early natural history of the eastern pygmy-possum, *Cercartetus nanus* (Geoffroy and Desmarest, 1817). *Proceedings of the Linnean Society of New South Wales* 127:107–124.
- HARRIS, J. M. 2008. *Cercartetus nanus* (Diprotodontia: Burramyidae). *Mammalian Species* 815:1–10.
- HARTMAN, C. G. 1940. Remarks on E. A. Bowley's note concerning alleged delayed fertilization in *Dromicia*. *Journal of Mammalogy* 21:213.
- HOLLAND, G. P. 1971. Two new species and three new subspecies of *Acanthopsylla* from Australia, with some notes on the genus (Siphonaptera: Pygiopsyllidae). *Canadian Entomologist* 103:213–234.
- HOPKINS, D., J. DICKSON, AND B. GAYNOR. 1984. Hepatoma in two pygmy possums (*Cercartetus concinnus*). *Australian Veterinary Journal* 61:265.
- HORNER, B. 1994. The importance of *Banksia marginata* and *Banksia ornata* as a food source for *Cercartetus concinnus* (the western pygmy-possum) and *Cercartetus lepidus* (the little pygmy-possum). Hons thesis, University of Adelaide, Adelaide, Australia (not seen, cited in New South Wales National Parks and Wildlife Service 2001).
- HOW, R. A., J. DELL, AND W. F. HUMPHREYS. 1987. The ground vertebrate fauna of coastal areas between Busselton and Albany, Western Australia. *Records of the Western Australian Museum* 13:553–574.
- HOW, R. A., J. DELL, AND B. G. MUIR. 1988. The biological survey of the eastern goldfields of Western Australia. Part 4. Lake Johnston–Hyden study area. IV. Vertebrate fauna. *Records of the Western Australian Museum Supplement* 30:44–83.
- INTERNATIONAL UNION FOR THE CONSERVATION OF NATURE AND NATURAL RESOURCES. 2006. 2006 IUCN Red list of threatened animals. www.iucnredlist.org, accessed 10 May 2008.
- IREDALE, T., AND E. L. G. TROUGHTON. 1934. A check-list of the mammals recorded from Australia. *Australian Museum Memoir* 6:1–122.
- JONES, A. D. 1954. The recent increase of the rarer native mammals. VII. Manjimup. *Western Australian Naturalist* 4:139–140.
- KAVANAGH, R. P. 2004. Distribution and conservation status of possums and gliders in New South Wales. Pp. 130–148 in *The biology of Australian possums and gliders* (R. L. Goldingay and S. M. Jackson, eds.). Surrey Beatty and Sons, Chipping Norton, Australia.
- KEAN, R. I. 1967. Marsupials (part 2). *Tuatara* 15:25–45.
- KEMP, L. F., AND S. M. CARTHEW. 2004. Nest site selection by the western pygmy-possum *Cercartetus concinnus*. Pp. 237–245 in *The biology of Australian possums and gliders* (R. L. Goldingay and S. M. Jackson, eds.). Surrey Beatty and Sons, Chipping Norton, Australia.
- KIRSCH, J. A. W. 1968. Prodomus of the comparative serology of Marsupialia. *Nature* 217:418–420.
- KITCHENER, D. J., A. CHAPMAN, B. G. MUIR, AND M. PALMER. 1980. The conservation value for mammals of reserves in the Western Australian wheatbelt. *Biological Conservation* 18:179–207.
- KITCHENER, D. J., AND E. VICKER. 1981. *Catalogue of modern mammals in the Western Australian Museum 1895 to 1981*. Western Australian Museum Publication, Perth, Australia.
- KREFFT, G. 1871. *The mammals of Australia, with a short account of all the species hitherto described*. Government Printer, Sydney, Australia.
- LANDWEHR, G. O., K. C. RICHARDSON, AND R. D. WOOLLER. 1990. Sugar preferences of honey possums *Tarsipes rostratus* (Marsupialia: Tarsipedidae) and western pygmy possums *Cercartetus concinnus* (Marsupialia: Burramyidae). *Australian Mammalogy* 13:5–10.
- LOARING, W. H. 1960. Pigmy possums at Bickley. *Western Australian Naturalist* 7(5):135.
- LOWRY, J. W. J., AND D. MERRILEES. 1969. Age of the desiccated carcass of a thylacine (Marsupialia, Dasyuroidea) from Thylacine Hole, Nullarbor Region, Western Australia. *Helictite* 7:15–16.
- LUNDELIUS, E. L., JR. 1957. Additions to the knowledge of the ranges of Western Australian mammals. *Western Australian Naturalist* 5:173–182.
- LUNDELIUS, E. L., JR. 1960. Post Pleistocene succession in Western Australia and its climatic interpretation. *Proceedings of the International Geological Congress* 21(4):142–153.
- LUNDELIUS, E. L., JR. 1963. Vertebrate remains from the Nullarbor caves, Western Australia. *Journal of the Royal Society of Western Australia* 46:75–80.
- LUNDELIUS, E. L., JR. 1983. Climatic implications of late Pleistocene and Holocene faunal associations in Australia. *Alcheringa* 7:125–149.
- LUNDELIUS, E. L., JR., AND W. D. TURNBULL. 1982. The mammalian fauna of Madura Cave, Western Australia, part V: Diprotodontia (part). *Fieldiana: Geology (New Series)* 11:1–32.
- LUNDELIUS, E. L., JR., AND W. D. TURNBULL. 1989. The mammalian fauna of Madura Cave, Western Australia, part VII: Macropodidae: Sthenurinae, Macropodinae, with a review of the marsupial portion of the fauna. *Fieldiana: Geology (New Series)* 17:1–71.
- LYDEKKER, R. 1896. *A handbook to the Marsupialia and Monotremata*. Llyod Natural History, London, United Kingdom.
- MARSHALL, A. J. 1967. Origin of delayed implantation in marsupials. *Nature* 216:192–193.

- MARTIN, P. G., AND D. L. HAYMAN. 1967. Quantitative comparisons between the karyotypes of Australian marsupials from three different superfamilies. *Chromosoma* 20:290–310.
- MATSCHIE, P. 1916. Die Verbreitung der Beuteltiere auf New Guinea mit einigen Bemerkungen über Ihre Einteilung in Untergattungen. *Mitteilungen Zoologisches Museum Berlin* 8:260.
- MCDOWELL, M. C. 1997. Taphonomy and palaeoenvironmental interpretation of a late Holocene deposit from Black's Point Sinkhole, Venus Bay, S.A. *Proceedings of the Linnean Society of New South Wales* 117:79–96.
- MCDOWELL, M. C. 2001. Analysis of late quaternary fossil mammal faunas from Robertson Cave (5U17, 18, 19) and Wet Cave (5U10, 11) in the Naracoorte World Heritage Area, South Australia. M.S. thesis, Flinders University, Adelaide, Australia.
- MCKAY, G. M. 1984. Cytogenetic relationships of possums and gliders. Pp. 9–16 in *Possums and gliders* (A. P. Smith and I. D. Hume, eds.). Surrey Beatty and Sons with Australian Mammal Society, Sydney, Australia.
- MCKAY, G. M. 1988. *Burramyidae*. Pp. 98–102 in *Zoological catalogue of Australia*. Vol. 5. Mammalia (J. L. Bannister, et al., eds.). Australian Government Publishing Service, Canberra, Australia.
- MCKENZIE, N. L., AND A. C. ROBINSON. 1987. A biological survey of the Nullarbor Region of South and Western Australia in 1984. South Australian Department of Environment and Planning, Western Australian Department of Conservation and Land Management, Australian National Parks and Wildlife Service, Canberra, Australia.
- MCKENZIE, N. L., AND J. K. ROLFE. 1995. The biological survey of the eastern goldfields of Western Australia. Part 11. Boorabbin–Southern Cross study area. IV. Vertebrate fauna. *Records of the Western Australian Museum Supplement* 49:31–65.
- MCKENZIE, N. L., J. K. ROLFE, N. J. HALL, AND W. K. YOUNGSON. 1983. The biological survey of the eastern goldfields of Western Australia. Part 9. Norseman–Balladonia study area. IV. Vertebrate fauna. *Records of the Western Australian Museum Supplement* 42:33–55.
- MENKHORST, P. W., AND C. M. BEARDSSELL. 1982. Mammals of southwestern Victoria from the Little Desert to the coast. *Proceedings of the Royal Society of Victoria* 94:221–247.
- MERRILEES, D. 1968. Man the destroyer: late Quaternary changes in the Australian marsupial fauna. *Journal of the Royal Society of Western Australia* 51:1–24.
- MERRILEES, D. 1984. Comings and goings of late Quaternary mammals in extreme southwestern Australia. Pp. 629–638 in *Quaternary extinctions: a prehistoric revolution* (P. S. Martin and R. G. Klein, eds.). University of Arizona Press, Tucson.
- MISSE, G. E. 1997. Nesting sites and habitat of the western pygmy-possum, *Cercartetus concinnus*. Hons thesis, University of Adelaide, Adelaide, Australia.
- MJÖBERG, E. 1916. On a new genus and species of marsupials. *Kungliga Svenska Vetenskapsakademiens Handlingar* 52(2):13–14.
- NEW SOUTH WALES NATIONAL PARKS AND WILDLIFE SERVICE. 2001. Western pygmy-possum (*Cercartetus concinnus*) approved recovery plan. New South Wales National Parks and Wildlife Service, Hurstville, New South Wales, Australia.
- NEW SOUTH WALES SCIENTIFIC COMMITTEE. 1997. Final determination to list the western pygmy-possum *Cercartetus concinnus* as an endangered species on Part 1 of Schedule 1 of the Threatened Species Conservation Act 1995. New South Wales Department of Environment and Climate Change, Sydney, Australia.
- Ogilby, J. D. 1892. Catalogue of Australian mammals, with introductory notes on general mammalogy. *Australian Museum Catalogue* 16:35.
- OSBORNE, M. J., AND L. CHRISTIDIS. 2002. Systematics and biogeography of pygmy possums (Burramyidae: *Cercartetus*). *Australian Journal of Zoology* 50:25–37.
- PARTRIDGE, J. 1967. A 3,300 year old thylacine (Marsupialia: Thylacynidae) from the Nullarbor Plain, Western Australia. *Journal of the Royal Society of Western Australia* 50:57–59.
- PATON, D. C. 1999. Impact of commercial honeybees on flora and fauna in Ngarkat Conservation Park. A report for the Rural Industries Research and Development Corporation, Barton, Australian Capital Territory, Australia.
- PEARSON, D. J., D. R. KING, AND E. R. PIANKA. 1999. New localities for the western pygmy possum, *Cercartetus concinnus*, in the Great Victoria Desert. *Western Australian Naturalist* 22:205–207.
- PESTELL, A. J. L., S. J. B. COOPER, K. M. SAINT, AND S. PETIT. 2008. Genetic structure of the western pygmy possum *Cercartetus concinnus* Gould (Marsupialia: Burramyidae) based on mitochondrial DNA. *Australian Mammalogy* 29:191–200.
- PESTELL, A. J. L., AND S. PETIT. 2007a. Methods and ethical considerations of pitfall trapping for the western pygmy possum (*Cercartetus concinnus* Gould) (Marsupialia: Burramyidae), with observations on capture patterns and nest sites. *Wildlife Research* 34:296–305.
- PESTELL, A. J. L., AND S. PETIT. 2007b. Diet of the western pygmy possum, *Cercartetus concinnus* Gould (Marsupialia: Burramyidae), at Innes National Park, South Australia, and evaluation of diet sampling methods. *Australian Journal of Zoology* 55: 275–284.
- PLEDGE, N. S. 1974. Fossil vertebrates. Pp. 123–127 in *Natural history of Kangaroo Island* (M. J. Tyler, G. R. Twidale, and J. K. Ling, eds.). Royal Society of South Australia, Adelaide, Australia.
- PORTER, J. K. 1979. Vertebrate remains from a stratified Holocene deposit in Skull Cave, Western Australia, and a review of their significance. *Journal of the Royal Society of Western Australia* 61: 109–117.
- REED, E. H., AND S. J. BOURNE. 2000. Pleistocene fossil vertebrate sites of the south east region of South Australia. *Transactions of the Royal Society of South Australia* 124:61–90.
- RENFREE, M. B. 1978. Embryonic diapause in mammals—a developmental strategy. Pp. 1–46 in *Dormancy and developmental arrest* (M. E. Cutter, ed.). Academic Press, New York.
- RENFREE, M. B. 1981. Embryonic diapause in marsupials. *Journal of Reproduction and Fertility Supplement* 29:67–78.
- RENFREE, M. B. 1993. Diapause, pregnancy, and parturition in Australian marsupials. *Journal of Experimental Zoology* 266: 450–462.
- RENFREE, M. B., AND J. H. CALABY. 1981. Background to delayed implantation and embryonic diapause. *Journal of Reproduction and Fertility Supplement* 29:1–9.
- RICHARDS, J. D., AND J. SHORT. 1996. History of the disappearance of native fauna from the Nullarbor Plain through the eyes of long time resident Amy Crocker. *Western Australian Naturalist* 21: 89–96.
- ROBERTSON, P., ET AL. 1989. Fauna of the mallee study area northwestern Victoria. Arthur Rylah Institute for Environmental Research Technical Report 87. Department of Conservation, Forests and Lands, Victoria, Australia.
- ROBINSON, A. C., AND D. M. ARMSTRONG (EDS.). 1999. A biological survey of Kangaroo Island, South Australia in November 1989 and 1990. Heritage and Biodiversity Section, Department for Environment, Heritage and Aboriginal Affairs, South Australia, Adelaide, Australia.
- ROFE, R., AND D. HAYMAN. 1985. G-banding evidence for a conserved complement in the Marsupialia. *Cytogenetics and Cell Genetics* 39:40–50.
- ROSE, A. 1995. The effects of rainfall on the abundance and species richness of small vertebrates in the Stirling Range National Park. *Western Australian Naturalist* 20:53–59.
- ROSENBERG, H. I., AND K. C. RICHARDSON. 1995. Cephalic morphology of the honey possum, *Tarsipes rostratus* (Marsupialia: Tarsipedidae); an obligate nectarivore. *Journal of Morphology* 223:303–323.
- ROSENBERG, H. I., AND R. ROSE. 1999. Volar adhesive pads of the feathertail glider, *Acrobates pygmaeus* (Marsupialia: Acrobatidae). *Canadian Journal of Zoology* 77:233–248.
- RUSSELL, E. M. 1982. Patterns of parental care and parental investment in marsupials. *Biological Reviews* 57:423–486.
- RYAN, R. M. 1963. Occurrence of the western pygmy possum, *Cercartetus concinnus*, in Victoria and New South Wales. *Victorian Naturalist* 79:337–340.
- SEEBECK, J., AND I. MANSERGH. 1998. Mammals introduced to Wilsons Promontory. *Victorian Naturalist* 115:350–356.
- SHARMAN, G. B. 1963. Delayed implantation in marsupials. Pp. 3–14 in *Delayed implantation* (A. C. Enders, ed.). University of Chicago Press, Chicago, Illinois.

- SHORTRIDGE, G. C. 1909. An account of the geographical distribution of the marsupials and monotremes of south-west Australia, having special reference to the specimens collected during the Balston Expedition of 1904–1907. *Proceedings of the Zoological Society of London* 55:803–848.
- SIMPSON, G. G. 1945. The principles of classification and a classification of mammals. *Bulletin American Museum Natural History* 85: 1–350.
- SINCLAIR KNIGHT MERZ. 2006. Report on fauna survey 2005: part I—Iluka Resources Ltd. Mineral Deposit Area, Yellabinnia Regional Reserve, South Australia. Sinclair Knight Merz, Hindmarsh, South Australia.
- SLAVEN, M. R., AND K. C. RICHARDSON. 1988. Aspects of the form and function of the kidney of the honey possum, *Tarsipes rostratus*. *Australian Journal of Zoology* 36:465–471.
- SMITH, M. J. 1984. Observations on the reproductive systems and paraoal glands of *Cercartetus lepidus* (Marsupialia: Burramyidae). *Australian Mammalogy* 7:175–178.
- SMITH, M. J. 1995. Western pygmy-possum *Cercartetus concinnus*. Pp. 213–214 in *The mammals of Australia* (R. Strahan, ed.). 2nd ed. Reed Books, Chatswood, Australia.
- STEWART, H. J., H. M. OWENS, G. CARPENTER, AND T. CROFT. 1998. A biological survey of Bunbury Conservation Reserve and Stoneleigh Park Heritage Agreement South Australia. Department for Environment, Heritage and Aboriginal Affairs, Adelaide, Australia.
- STRAHAN, R. 1981. A dictionary of Australian mammal names: pronunciation, derivation, and significance of the names; with biographical and bibliographical notes. Angus and Robertson Publishers, Sydney, Australia.
- THOMAS, O. 1888. Catalogue of the Marsupialia and Monotremata in the collections of the British Museum (Natural History). British Museum (Natural History) Publications, London, United Kingdom.
- THOMAS, O. 1922. A selection of lectotypes of the typical Australian marsupials in the British Museum collection. *Annals and Magazine of Natural History* 9(10):127–128.
- TIDEMANN, C. R. 1967. Some mammal remains from cave deposits in the south-east of South Australia. *South Australian Naturalist* 42: 21–26.
- TULLOCH, A. 2004. The importance of food and shelter for habitat use and conservation of the burramyids in Australia. Pp. 268–284 in *The biology of Australian possums and gliders* (R. L. Goldingay and S. M. Jackson, eds.). Surrey Beatty and Sons, Chipping Norton, Australia.
- TUNBRIDGE, D. 1991. The story of the Flinders Ranges mammals. Kangaroo Press, Kenthurst, Australia.
- TURNER, V. 1982. Marsupials as pollinators in Australia. Pp. 55–66 in *Pollination and evolution* (J. A. Armstrong, J. M. Powell, and A. J. Richards, eds.). Royal Botanic Gardens Trust, Sydney, Australia.
- TURNER, V., AND G. M. MCKAY. 1989. Burramyidae. Pp. 652–664 in *Fauna of Australia*. Vol. 1B Mammalia (D. W. Walton and B. J. Richardson, eds.). Australian Government Publishing Service, Canberra, Australia.
- TYNDALE-BISCOE, C. H. 1973. *Life of marsupials*. Arnold, London, United Kingdom.
- TYNDALE-BISCOE, H., AND M. RENFREE. 1987. *Reproductive physiology of marsupials*. Cambridge University Press, Melbourne, Australia.
- VAN DER REE, R., S. J. WARD, AND K. A. HANDASYDE. 2004. Distribution and conservation status of possums and gliders in Victoria. Pp. 91–110 in *The biology of Australian possums and gliders* (R. L. Goldingay and S. M. Jackson, eds.). Surrey Beatty and Sons, Chipping Norton, Australia.
- VAN DYCK, S., AND R. STRAHAN (EDS.). 2008. *The mammals of Australia*. 3rd ed. Reed New Holland, Sydney, Australia.
- WAITE, E. R., AND F. WOOD JONES. 1927. The fauna of Kangaroo Island, South Australia. 2. Mammals. *Transactions and Proceedings of the Royal Society of South Australia* 51:322–325.
- WAKEFIELD, N. A. 1963. The Australian pigmy possums. *Victorian Naturalist* 80:99–116.
- WAKEFIELD, N. A. 1966. Mammals recorded for the mallee, Victoria. *Proceedings of the Royal Society of Victoria* 79:627–633.
- WAKEFIELD, N. A. 1970. Notes on Australia pigmy-possums (*Cercartetus*, Phalangeridae, Marsupialia). *Victorian Naturalist* 87:11–18.
- WAKEFIELD, N. A. 1971. Mammals of western Victoria. Pp. 35–51 in *The natural history of western Victoria* (M. H. Douglas and L. O'Brien, eds.). Australian Institute of Agricultural Science, Horsham, Victoria, Australia.
- WARD, S. J. 1990. Reproduction in the western pygmy-possum, *Cercartetus concinnus* (Marsupialia, Burramyidae), with notes on reproduction of some other small possum species. *Australian Journal of Zoology* 38:423–438.
- WARD, S. J. 1992. Life history of the little pygmy-possum, *Cercartetus lepidus* (Marsupialia: Burramyidae), in the Big Desert, Victoria. *Australian Journal of Zoology* 40:43–55.
- WARD, S. J. 1998. Numbers of teats and pre- and post-natal litter sizes in small diprotodont marsupials. *Journal of Mammalogy* 79:999–1008.
- WATERHOUSE, G. R. 1846. *A natural history of the Mammalia*. Vol. 1. Containing the order Marsupialia, or pouched animals. Baillière, London, United Kingdom.
- WATTS, C. H. S. 1990. A list of the vertebrates of South Australia. Department of Environment and Planning, Adelaide, South Australia.
- WATTS, C. H. S., AND J. K. LING. 1985. Marine and terrestrial mammals. Pp. 139–147 in *Natural history of Eyre Peninsula* (C. R. Twidale, M. J. Tyler, and M. Davies, eds.). Royal Society of South Australia, Northfield, Australia.
- WESCOTT, G. 1998. The scientific importance of Wilsons Promontory. *Victorian Naturalist* 115:296–299.
- WHITTELL, H. M. 1954. John Gilbert's notebook on marsupials. *Western Australian Naturalist* 4(5):104–114.
- WILLIAMS, D. L. G. 1980. Catalogue of Pleistocene vertebrate fossils and sites in South Australia. *Transactions of the Royal Society of South Australia* 104:101–115.
- WOOD JONES, F. 1924. *The mammals of South Australia*. Part II. Government Printer, Adelaide, Australia.
- WOOD JONES, F. 1925. A new South Australian dormouse opossum. *Transactions of the Royal Society of South Australia* 49:96–98.
- WOOLLER, R. D., K. C. RICHARDSON, AND G. O. BRADLEY. 1999. Dietary constraints upon reproduction in an obligate pollen- and nectar-feeding marsupial, the honey possum (*Tarsipes rostratus*). *Journal of Zoology (London)* 248:279–287.
- WOOLLEY, P. A., AND A. E. VANDERVEEN. 2002. The rete testis of acrobatid and burramyid marsupials. *Australian Journal of Zoology* 50:237–247.
- ZOSKY, G. R., AND A. N. LARCOMBE. 2003. The parasymphathetic nervous system and its influence on heart rate in torpid western pygmy possums, *Cercartetus concinnus* (Marsupialia: Burramyidae). *Zoology* 106:143–150.

Associate editors of this account were PAMELA OWEN and DAVID ZEGERS. COLIN GROVES reviewed the synonymies. Editor was MEREDITH J. HAMILTON.