

## Peromyscus eva.

By Sergio Ticul Álvarez-Castañeda and Patricia Cortés-Calva

Published 18 December 2003 by the American Society of Mammalogists

### Peromyscus eva Thomas, 1898

#### Baja California Sur Deer Mouse

*Peromyscus eva* Thomas, 1898:44. Type locality "San José del Cabo, Baja California."

*Peromyscus eremicus propinquus* Allen, 1898:154. Type locality "San Pablo point, Baja California."

*Peromyscus eremicus carmeni* Townsend, 1912:154. Type locality "Carmen island [Isla del Carmen], Gulf of California, Baja California."

**CONTEXT AND CONTENT.** Order Rodentia, suborder Sciurognathii, family Muridae, subfamily Sigmodontinae, genus *Peromyscus* (Mussner and Carleton 1993), subgenus *Haplomylomys*. Two subspecies are recognized currently (Álvarez-Castañeda and Cortés-Calva 1999).

*P. e. carmeni* Townsend, 1912:154, see above.

*P. e. eva* Thomas, 1898:44, see above.

**DIAGNOSIS.** *Peromyscus eva* (Fig. 1) has a gray face and a short pelage with a variable blend of russet, buff, and brown with finely textured appearance (Lawlor 1971). *P. eva* and *P. eremicus* (*P. fraterculus* sensus Hafner et al. 2001) have more or less the same colors, but *P. fraterculus* is noted for a mixture of dark brown or black (Lawlor 1971). Tail of *P. eva* is longer and ears are slightly smaller than those of *P. fraterculus* (= *P. eremicus*—Osgood 1909). In the Isla del Carmen population, *P. eva* is darker and similar to mainland *P. fraterculus*. Compared to *P. fraterculus*, *P. eva* is has larger tail length, greatest length of skull, zygomatic breath, zygomatic notch, naso frontal suture forward of the maxillary-frontal suture, and length and width of toothrows. Bacular morphology differentiates *P. eva* and *P. fraterculus* (Burt 1960 in Lawlor 1971).

**GENERAL CHARACTERISTICS.** Dorsal surface of *P. eva* is ocherous buff, mixed conspicuously with uniformly distributed fine dusky dorsal lines, except on lower sides. An ocherous buff lateral line, usually distinct only from axillary region to hip, is often wider medially. Head, including nose, cheeks, and orbital region, is pale gray tinged with ocherous buff. A narrow dusky orbital ring extends scarcely beyond eyelids. Ears are pale brown and almost hairless. Ventrum is pure creamy white without pectoral spot. Feet are white (Osgood 1909). Zygoma is generally rounded laterally, and vertebrae number <36 (Fig. 2; Lawlor 1971).

Average measurements with parenthetical SD and range for 8 female and 10 male *P. e. carmeni* from Isla del Carmen, México, in the Centro de Investigaciones Biológicas del Noroeste collection are (mm): length of head and body,  $185.3 \pm 6.2$  (175–197); length of tail,  $104.5 \pm 4.2$  (97–113); length of hind foot,  $19.3 \pm 1.1$  (17–21); length of ear,  $18.1 \pm 1.8$  (15–22); greatest length of skull,  $24.3 \pm 0.6$  (22.9–25.1); palatilar length,  $11.2 \pm 0.3$  (10.7–11.7); cranial breadth,  $11.6 \pm 0.3$  (11.1–12.0); interorbital breadth,  $4.0 \pm 0.1$  (3.8–4.4); bullar length,  $3.1 \pm 0.2$  (2.8–3.6); mandibular length,  $11.9 \pm 0.3$  (11.1–12.5); length of maxillary toothrow,  $3.6 \pm 0.1$  (3.3–3.8); length of mandibular toothrow,  $3.9 \pm 0.2$  (3.4–4.2). Mean mass is  $15.5 \pm 1.5$  g (12.7–17.5).

**DISTRIBUTION.** *Peromyscus eva* occurs on the Baja California Peninsula from the cape region to the border area between the states of Baja California and Baja California Sur (Carleton 1980), including Isla del Carmen (Lawlor et al. 2002). In its southern range, it is not present above 1,800 m in the Sierra de la Laguna (Banks 1967a), nor in the northern part of its range, above 1,200 m in the Sierra de la Giganta (Álvarez 1958). Only 1 specimen was previously recorded, from Isla del Carmen off the coast at Loreto, Baja California Sur (Huey 1964; Lawlor 1983). No fossils are known (Hibbard 1968).

**FORM AND FUNCTION.** *Peromyscus eva* changes considerably over its geographic range. Specimens in the southern area are smaller than those in the north (Lawlor 1971). Color of dorsum varies from bright, sandy russet in the south to ocherous brown in the north (Lawlor 1971). Dorsal hair of the Isla del Carmen population is gray with ocherous tips, and sides are more ocherous than dorsum.

Average bacula measurements (mm) of 3 specimens from Cabo San Lucas are length 8.1 (7.9–8.3) and width 0.8 (0.8–0.8). One baculum from Isla del Carmen was 8.0 mm long and 0.7 wide (Burt 1960). Dental formula is i 1/1, c 0/0, p 0/0, m 3/3, total 16.

**REPRODUCTION.** Females of *P. eva* captured near La Paz during February, March, and June were pregnant, and lactation was observed mainly in February, April, and July. Males had scrotal testes mainly from December to May. Breeding was observed in July (Cortés-Calva in litt.).

In May, 1 male *P. e. carmeni* had 5.9-mm testes. In July, 3 pregnant and lactating females were captured. Two embryos measured 7.7 by 5 mm. Number of embryos averaged 3, and size of embryos ranged from 1.5–22 mm (Sánchez et al. 1997). Testes of 2 mainland males in May were 4 and 5 mm, and those of 1 male in June were 5 mm (Sánchez et al. 1997).

**ECOLOGY.** *Peromyscus eva* was collected in fields of chili, corn, oranges, palms, sugarcane, and tamarind; in areas with secondary vegetation; and in natural arid tropical scrub (Álvarez-Castañeda 1994). It occurred in stony areas and natural habitats containing matorral sarcocaula such as chain fruit cholla/cholla (*Cylindropuntia opuntia*), Adam's tree/palo Adam (*Fouquieria diguetii*), ashy limberbush/lomboy (*Jatropha cinerea*), viejito (*Mammillaria*), cardon (*Pachicereus pringlei*), and organ-pipe cactus/pitaya agria (*Stenocereus gummosus*—Cortés-Calva and Álvarez-Castañeda 1997). The population of *P. eva* near La Paz increases more during the cooler months October to February than other species of rodents, such as *Chaetodipus arenarius*, *C. ruddinoris*, *C. spinatus*, and *Dipodomys merriami*. *P. eva* is associated with patches of succulent vegetation (Cortés-Calva, in litt.).

On the slopes of the Sierra de la Laguna, the relative abundances of *P. eva* and *P. truei* are closely related to elevation and vegetation (Banks 1967a; Woloszyn and Woloszyn 1982). *P. eva*



FIG. 1. Subadult male *Peromyscus eva eva* from north of La Paz, Baja California Sur, Mexico. Photograph by Patricia Cortés-Calva.




FIG. 2. Dorsal, ventral, and lateral views of cranium and lateral view of mandible of *Peromyscus eva carmeni* (subadult, female, Isla del Carmen, Baja California Sur, Mexico, in the collection of the Centro de Investigaciones Biológicas del Noroeste, number 567). Greatest length of cranium is 24.8 mm. Photograph by A. Gutiérrez.

and *P. fraterculus* are sympatric in central Baja California Peninsula (Lawlor 1971).

*Peromyscus eva carmeni* is endemic to Isla del Carmen, 15 km east of Loreto. The closest distance from this island to the peninsula is 6 km (Nieto-Garibay 1999). The island measures 27 by 9 km (151 km<sup>2</sup>). A range of altitude 479 m crosses the island. *P. e. carmeni* occupies the stony areas of valleys and slopes (Sánchez et al. 1997). Dominant vegetation includes elephant tree/torote (*Bursera microphylla hindsiana*), desmanthus/frijolillo (*Desmanthus fruticosus*), spurge/liga (*Euphorbia polycarpa*), barrel cactus/viznaga (*Ferocactus diguetii*, var. *carmensis*), Adam's tree/palo Adan (*Fouquieria diguetii*), leatherplant/matacora (*Jatropha curneata*), ironwood/palo fierro (*Olnaea tesota*), chain fruit cholla/cholla (*Opuntia burrageana*), cardon (*Pachicereus pringlei*), Spanish needle/aguja (*Palaeoxia leucophylla*), passion flower/granadilla (*Pasiflora palmeri*), slipper plant/candelilla (*Pedilanthus macrocarpus*), honey mesquite/mesquite (*Prosopis glandulosa*), pickleweed/salicornia (*Salicornia subterminalis*), and organ-pipe cactus/pitaya agria (*Stenocereus gummosus*—Moran 1983).

*Peromyscus eva* hosts the following parasites: Acarina: *Hex-*




FIG. 3. Distribution of *Peromyscus eva* in Baja California Sur, Mexico (modified from Hafner et al. 2001): 1, *P. e. carmeni*; 2, *P. e. eva*.

*idionis macropus* (Loomis and Lucas 1970) and Coccidia: *Eimeria arizonensis* and *E. langebarteli* (McAllister et al. 1993).

*P. eva* is prey to *Lanius ludovicianus* (Álvarez-Castañeda 2002), *Bubo virginianus* (Llinás-Gutiérrez et al. 1991), *Tyto alba* (López-Forment and Urbano 1977), *Canis latrans* (López-Forment et al. 1996), *Urocyon cinereoargenteus* (Arnaud and Acevedo 1990), and snakes (Townsend 1912). Other mammals associated with *P. eva* are *Ammospermophilus leucurus*, *Chaetodipus arenarius*, *C. rufinoris*, *C. spinatus*, *Dipodomys merriami*, *Lepus californicus*, *Neotoma lepida*, *Peromyscus fraterculus*, *P. maniculatus*, *P. truei*, and *Sylvilagus audubonii* (Álvarez-Castañeda 1994; Banks 1967b; Cortés-Calva and Álvarez-Castañeda 1997).

**GENETICS.** Karyotype of *P. eva* closely resembles that of *P. eremicus*. All chromosomes are biaxmed,  $2n = 48$  (Lawlor 1971). Allozymes of *P. e. carmeni* differ from *P. eremicus* of the Baja California Peninsula (Avise et al. 1974).

**CONSERVATION STATUS.** *P. eva carmeni* is listed as threatened in the Norma Oficial Mexicana (2001). Factors that threaten *P. e. carmeni* in Isla del Carmen include nonnative species such as cats and goats and human activities like fishing, mining, and tourism (Álvarez-Castañeda and Ortega-Rubio 2003). *P. e. eva* does not have conservation status in Mexico.

**REMARKS.** *P. eva* was described as a species by Thomas (1898) but was considered a subspecies of *P. eremicus* by Osgood (1909) and Hooper (1968). Burt (1960) used bacular differences to distinguish *P. eva* from *P. eremicus*. Lawlor (1971) mentioned that *P. eva* was derived from an *eremicus*-like ancestor and said that some specimens of the Sierra de la Giganta have features that are combinations of *P. eremicus* and *P. eva*. Molecular techniques suggest a sister-group relationship between *eva* and *eremicus*-like forms on the Baja Peninsula (Riddle et al. 2000). The Latin name *carmeni* derives from the Isla del Carmen.

We thank the editing staff at CIBNOR for editing the text. Financial support was provided by Consejo Nacional de Ciencia y Tecnología (CONACYT grant J28319-N).

#### LITERATURE CITED

- ÁLVAREZ, T. 1958. Roedores colectados en el territorio de la Baja California. Acta Zoológica Mexicana 2:1–7.
- ÁLVAREZ-CASTAÑEDA, S. T. 1994. Current status of the rice rat, *Oryzomys couesi peninsularis*. Southwestern Naturalist 39: 99–100.
- ÁLVAREZ-CASTAÑEDA, S. T. 2002. Predation of desert mammals

- by *Lanius ludovicianus* (Laniidae). Southwestern Naturalist 7: 628–630.
- ÁLVAREZ-CASTAÑEDA, S. T., AND P. CORTÉS-CALVA. 1999. Familia Muridae. Pp. 445–566 in Mamíferos del noroeste de México (S. T. Álvarez-Castañeda and J. L. Patton, eds.). Centro de Investigaciones Biológicas del Noroeste, La Paz, B.C.S., México.
- ÁLVAREZ-CASTAÑEDA, S. T., AND A. ORTEGA-RUBIO. 2003. Current status of rodents on islands in the Gulf of California. Biological Conservation 109:157–163.
- ARNAUD, G., AND M. ACEVEDO. 1990. Hábitos alimenticios de la zorra gris, *Urocyon cinereoargenteus* (Carnívora: Canidae) en la región meridional de Baja California, México. Revista de Biología Tropical 32:499–502.
- AVISE, J. C., M. H. SMITH, R. K. SELANDER, T. E. LAWLER, AND P. R. RAMSEY. 1974. Biochemical polymorphism and systematic in the genus *Peromyscus*. V. Insular and mainland species of the subgenus *Haplomylomys*. Systematic Zoology 23:226–238.
- BANKS, R. C. 1967a. Birds and mammals of La Laguna, Baja California. Transactions of the San Diego Society of Natural History 14:205–232.
- BANKS, R. C. 1967b. Birds and mammals of the voyager of the “Gringa”. Transactions of the San Diego Society of Natural History 13:177–184.
- BURT, W. H. 1960. Catalog of North American mammals. Miscellaneous Publications, Museum of Zoology, University of Michigan 113:1–76.
- CARLETON, M. D. 1980. Systematic and evolution. Pp. 7–141 in Advances in the study of *Peromyscus* (Rodentia) (G. L. Kirkland Jr. and J. N. Layne, eds.). Texas Tech University Press, Lubbock.
- CORTÉS-CALVA, P. In press. Estudio de la comunidad de roedores relacionados a matorral árido tropical de Baja California Sur, México. Ph.D. dissertation, Facultad de Ciencias, Universidad Nacional Autónoma de México.
- CORTÉS-CALVA, P., AND S. T. ÁLVAREZ-CASTAÑEDA. 1997. Diversidad de roedores en zona de la bahía de La Paz. Pp. 265–272 in La Bahía de La Paz, investigación y conservación (J. Urban R. and M. Ramírez R., eds.). Universidad Autónoma de Baja California Sur and Scripps Institution of Oceanography, La Paz, B.C.S., and La Jolla, California.
- HAFNER, D. J., B. R. RIDDLE, AND S. T. ÁLVAREZ-CASTAÑEDA. 2001. Evolutionary relationships of white-footed mice (*Peromyscus*) on islands in the Sea of Cortez, Mexico. Journal of Mammalogy 82:775–790.
- HIBBARD, C. W. 1968. Paleontology. Pp. 6–26 in Biology of *Peromyscus* (Rodentia) (J. A. King, ed.). Special Publication, American Society of Mammalogists 2:1–593.
- HOOPER, E. T. 1968. Classification. Pp. 27–74 in Biology of *Peromyscus* (Rodentia) (J. A. King, ed.). Special Publication, American Society of Mammalogy 2:1–593.
- HUEY, L. M. 1964. The mammals of Baja California, México. Transactions of the San Diego Society of Natural History 13: 85–168.
- LAWLER, T. E. 1971. Distribution and relationships of six species of *Peromyscus* in Baja California and Sonora, Mexico. Occasional Papers of the Museum of Zoology, University of Michigan 661:1–22.
- LAWLER, T. E. 1983. The mammals. Pp. 265–289 in Island biogeography in the Sea of Cortez (T. J. Case and M. L. Cody, eds.). University of California Press, Berkeley.
- LAWLER, T. E., D. J. HAFNER, B. R. RIDDLE, S. T. ÁLVAREZ-CASTAÑEDA, AND P. STAPP. 2002. The mammals. Pp. 326–361 in A new island biogeography of the Sea of Cortéz. Second edition (T. J. Case, M. L. Cody, and E. Ezcurra, eds.). Oxford University Press, United Kingdom.
- LLINAS-GUTIERREZ, J., G. ARNAUD, AND M. ACEVEDO. 1991. Food habits of the great horned owl (*Bubo virginianus*) in the Cape region of Lower California, México. Journal of Raptor Research 25:140–141.
- LOOMIS, R. B., AND J. L. LUCAS. 1970. A new subgenus and two new species of *Hexidionis* (Acarina, Tronbiculidae) from North America. Bulletin of the South California Academy of Sciences 69:52–59.
- LÓPEZ-FORMENT, W., I. E. LIRA, AND C. MUDESPACHER. 1996. Mamíferos: su diversidad en las islas mexicanas. AGT editores, S.A., México City, México.
- LÓPEZ-FORMENT, W., AND G. URBANO. 1977. Restos de pequeños mamíferos recuperados en regurgitaciones de lechuza, *Tyto alba*, en México. Anales del Instituto de Biología, Universidad Nacional Autónoma de México 48:231–242.
- MORAN, R. 1983. Vascular Plants of the Gulf Islands. Pp. 348–381 in Island biogeography in the Sea of Cortez (T. J. Case and M. L. Cody, eds.). University of California Press, Berkeley.
- MCAULIFFER, C. T., S. J. UPTON, R. R. HOLLANDER, AND K. M. HOGAN. 1993. Coccidian parasites of heteromyid and murid rodents from Baja California del Sur, Mexico. Journal of Helminthology Society of Washington 60:256–258.
- MUSSER, G. G., AND M. D. CARLETON. 1993. Family Muridae. Pp. 501–756 in Mammals species of the world, a taxonomic and geographic reference. Second edition (D. E. Wilson and D. M. Reeder, eds.). Smithsonian Institution Press, Washington, D.C.
- NIETO-GARIBAY, A. 1999. Características generales del Noroeste de México. Pp. 13–28 in Mamíferos del noroeste de México (S. T. Álvarez-Castañeda and J. L. Patton, eds.). Centro de Investigaciones Biológicas del Noroeste, La Paz, B.C.S. México.
- NORMA OFICIAL MEXICANA. 2001. Proyecto de Norma Oficial Mexicana Proy-NOM-059-Ecol-2000. Protección Ambiental. Especies de flora y fauna silvestres de México. Categorías de riesgo y especificaciones para su inclusión, exclusión o cambio. Lista de especies en riesgo. Diario oficial de la federación. 16 de Octubre, Mexico City, Mexico.
- OSSGOOD, W. H. 1909. Revision of the mice of the American genus *Peromyscus*. North American Fauna 28:1–268.
- RIDDLE, B. R., D. J. HAFNER, AND L. F. ALEXANDER. 2000. Phylogeography and systematics of the *Peromyscus eremicus* species-group on the historical biogeography of North America warm regional deserts. Molecular Phylogenetics and Evolution 17:145–160.
- SÁNCHEZ H., C., S. SANTILLAN A., AND I. ANAYA C. 1997. Observaciones sobre mamíferos terrestres de algunas islas del Golfo de California. Pp. 325–342 in Homenaje al profesor Ticul Álvarez (J. Arroyo C. and O. J. Polaco, eds.). Colección científica, Instituto Nacional de Antropología e Historia, Mexico, D.F.
- THOMAS, O. 1893. On a new mammals from western Mexico and Lower California. Annals Magazine of Natural History, series 7, 1:40–46.
- TOWNSEND, C. H. 1912. Mammals collected in Lower California, with descriptions of new species. American Museum of Natural History 31:117–130.
- WOLOSZYN, D., AND B. W. WOLOSZYN. 1982. Los mamíferos de la Sierra de la Laguna Baja California Sur. Consejo Nacional de Ciencia y Tecnología, México, D.F.

Associate editor of this account was LUI MARINELLI. Editor was VIRGINIA HAYSEN.

SÉRGIO TICUL ÁLVAREZ-CASTAÑEDA AND PATRICIA CORTÉS-CALVA, CENTRO DE INVESTIGACIONES BIOLÓGICAS DEL NOROESTE, APARTADO POSTAL 128, LA PAZ, BAJA CALIFORNIA SUR, C.P. 23000, MÉXICO.